arXiv:hep-th/9805085v1 14 May 1998

UWThPh-22-1998

February 7, 2008

A Treatment of the Schwinger Model within Noncommutative Geometry

H. Grosse

Institut for Theoretical Physics, University of Vienna,

Boltzmanngasse 5, A-1090 Vienna, Austria

P. Prešnajder[1]

Department of Theoretical Physics, Comenius University

Mlynská dolina, SK-84215 Bratislava, Slovakia

Abstract

We describe a free spinor field on a noncommutative sphere starting from a canonical realization of the algebra $\mathcal{U}(u(2|1))$ and a sequence of $su(2)$-invariant embeddings of $su(2)$ representations. The gauge extension of the model - the Schwinger model on a noncommutative sphere is defined and the model is quantized. Due to the noncommutativity of the sphere, the model contains only a finite number of modes, and consequently is non-perturbatively UV-regular. The fermionic determinants and the effective actions are calculated. The origin of chiral anomaly is clarified. In the commutative limit standard formulas are recovered.

---

[1] Supported by the ”Fond zur Förderung der Wissenschaftlichen Forschung in Österreich” project P11783-PHY and by the Slovak Grant Agency VEGA project 1/4305/97.

1 Introduction

The basic notions of the noncommutative geometry were developed in [1], [2], [3], and in the form of the matrix geometry in [4], [5]. The essence of this approach consists in reformulating first the geometry in terms of commutative algebras and modules of smooth functions, and then generalizing them to their noncommutative analogs. The notion of the space as a continuum of points is lost, and this is expected to lead to an UV-regular quantum field theory.

One of the simplest models for a noncommutative manifold is the noncommutative (fuzzy) sphere. It was introduced by many authors using various techniques, [6], [7], [8], [9]. In general, these are related to finite dimensional representations of the $SU(2)$ group. Thus, the models in question are basically matrix models. Quantum field theoretical models with a self-interacting scalar fields on a truncated sphere were described in [10], [11]. Since, the fields posses only finite number of modes the models are UV-regular.

The basic ingredient of the noncommutative geometry developed in [1], [2] is the spectral triple $(\mathcal{A}, D, \mathcal{H})$, together with a chirality operator (grading Γ) and a charge conjugation (antilinear isometry $\mathcal{J}$). Besides a noncommutative algebra $\mathcal{A}$, the spectral consists of a Dirac operator D acting in the Hilbert space of spinor fields $\mathcal{H}$. The free spinor fields on a noncommutative sphere in the spirit of noncommutative geometry were introduced in [11] and [12] in the framework of supersymmetric approach. The spectrum of the free Dirac operator was found identical but truncated to the standard one on a commutative sphere.

Our aim is to demonstrate, that the axioms of the noncommutative geometry can be implemented to a nontrivial field theoretical model. The model we wish to describe is the noncommutative analog of the Schwinger model on a sphere. The commutative quantum version was analyzed in detail in [14]. Its noncommutative matrix version was proposed in [15], an approach going behind matrix models was sketched in [16]. Recently a systematic classical noncommutative formulation was found in [17], where a differential calculus on a supersphere in the commutative and noncommutative cases is described in detail. Gauge models (for topologically trivial field configurations) are described in both commutative and noncommutative versions.

Here we shall first modify the supersymmetric descpription of topologically nontrivial spinor field configurations on a noncommutative sphere found in [11]. Instead of using in the formulation of the model finite dimensional representations of the superalgebra $osp(2|1)$ (thus working in fact within the matrix model approach), we shall start from an infinite dimensional canonical realization of the enveloping algebra $\mathcal{U}(u(2|1))$. We introduce linear $su(2)$-invariant embeddings of various sequences of finite dimensional representations of the even $su(2)$ subalgebra (similar embeddings are important within a general approach to the quantization of vector bundles proposed in [18]; this is natural since, our interpretation of fields on a nocommutative sphere is technically close to the mentioned quantization). Only the evaluation of the action is performed within particular finite dimensional representations. A free Dirac operator is introduced and the model is gauged. After specification of the gauge degrees of freedom the model is quantized. The resulting model is well defined nonperturbatively and UV-regular. This allows an exact analysis of various non-perturbative objects like fermionic determinants and effective actions. The non-perturbative origin of chiral symmetry breaking is clarified. In the commutative limit the standard formulas are reproduced (see e.g. [14]).

The paper is organized as follows. In Section 2 we describe the free spinor field on a noncommutative sphere within the $u(2|1)$ supersymmetric formalism and we discuss linear embeddings of finite dimensional representations of the even $su(2)$ subalgebra. In Section 3 we introduce gauge degrees of freedom, and we present a field action for noncommutative Schwinger model. In Section 4 we quantize nonperturbatively the model, and then we calculate fermionic determinants and effective field actions. Last Section 5 contains concluding remarks.

2 Free spinor field

First we summarize the commutative version of the model in question in the $SU(2)$-invariant supersymmetric formulation (see [14] for $SU(2)$-invariant formalism, and [11], [12], [17] for its supersymmetric reformulation). In this approach the fields are functions of two pairs χ_α, χ^*_α, $\alpha = 1, 2$, of complex variables and of one anticommuting (Grassmannian) pair a, a^*:

$$\Phi \ = \ \sum a^{\mu\nu}_{mn} \chi^{*m} \chi^n a^{*\mu} a^\nu \ , \tag{1}$$

where $n = (n_1, n_2)$ is a two component index and $\mu, \nu = 0, 1$ (we are using a multiindex notations: $\chi^n = \chi^{n_1}, \chi^{n_2}$, $|n| = n_1 + n_2$, $n! = n_1!n_2!$, etc.). The space $s\mathcal{H}_k$ of fields with a topological winding number $2k$ is defined as the space of fields (1) with $2k = |n| - |m| + \mu - \nu$ fixed. Any field from $s\mathcal{H}_k$ can be expanded as

$$\Phi \ = \ \Phi_0(\chi,\chi^*) \ + \ f(\chi,\chi^*)a \ + \ g(\chi,\chi^*)a^* \ + \ F(\chi,\chi^*)\gamma \ , \tag{2}$$

where $\gamma = a^*a - aa^*$, $\Phi_0, F \in \mathcal{H}_k$, $f \in \mathcal{H}_{k+\frac{1}{2}}$ and $g \in \mathcal{H}_{k-\frac{1}{2}}$. Here $\mathcal{H}_{k'}$ denotes the subspace of fields from $s\mathcal{H}_{k'}$ with $\mu = \nu = 0$.

The subspace of odd elements from $s\mathcal{H}_k$ is identified with the space $\mathcal{S}_k$ of spinor fields with a given topological winding number $2k$, i.e. any field from $\mathcal{S}_k$ can be expanded as

$$\Psi \ = \ f(\chi,\chi^*)a \ + \ g(\chi,\chi^*)a^* \ , \tag{3}$$

where $f \in \mathcal{H}_{k+\frac{1}{2}}$ and $g \in \mathcal{H}_{k-\frac{1}{2}}$. The chirality operator Γ in $\mathcal{S}_k$ is given as $\Gamma = P_+ - P_-$, where $P_\pm$ are projectors $\mathcal{S}_k \to \mathcal{S}_k^\pm$ given by:

$$P_+\Psi \ = \ a\partial_a\Psi \ = \ fa \ , \ P_-\Psi \ = \ a^*\partial_{a^*}\Psi \ = \ ga^* \ . \tag{4}$$

The chirality operator takes in $\mathcal{S}_k^\pm$ the value ± 1. The charge conjugation $\mathcal{J}$ is defined as follows

$$\mathcal{J}\Psi \ = \ \mathcal{J}(fa \ + \ ga^*) \ := \ g^*a \ - \ f^*a^* \ = \ \Gamma\Psi^* \ = \ \bar{\Psi} \ . \tag{5}$$

Obviously, $\mathcal{J} : \mathcal{S}_k \to \mathcal{S}_{-k}$, and $\mathcal{J}^2 = -1$. The inner product in $\mathcal{S}_k$ we define as

$$(\Psi_1, \Psi_2) \ = \ \int d\mu \bar{\Psi}_1 \Psi_2 \ = \ \int d\nu (f_1^* f_2 + g_1^* g_2) \ , \tag{6}$$

where $d\mu = \frac{1}{16\pi^2 r} d^2\chi d^2\chi^* da da^* \delta(\chi_\alpha^*\chi_\alpha + a^*a - r)$ and $d\nu = \frac{1}{2\pi r} d^3x \delta(x_i^2 - r^2)$ are normalized measures on a supersphere sS^3 and sphere S^2 respectively.

In the superspace $\mathbf{C}^{2|1}$ we have a natural action of the superalgebra $u(2|1)$ of matrices [19]

$$\begin{pmatrix} X & 0 \\ 0 & B \end{pmatrix} + \frac{1-i}{\sqrt{2}} \begin{pmatrix} 0 & V \\ -V^* & 0 \end{pmatrix} \ , \ X \ = \ -X^* \ , \ B \ = \ -B^* \ , \tag{7}$$

written in a (2|1)-block diagonal form. In the space $\mathbf{C}^{2|1}$ a super-Poisson bracket algebra can be introduced by postulating elementary bracket relations among $\mathbf{C}^{2|1}$ coordinate functions

$$[\chi_\alpha, \chi^*_\beta] \ = \ \delta_{\alpha\beta} \ , \ [a, a^*] \ = \ 1 \tag{8}$$

(with all other elementary brackets vanishing). The $u(2|1)$ superalgebra is then realized in terms of super-Poisson brackets by choosing its basis in the following way:

$$\begin{aligned} x_i \ &= \ \frac{i}{2}\chi^+\sigma_i\chi \ , \ b \ = \ \chi^+\chi \ + \ 2a^*a \ , \\ v_\alpha \ = \ \chi_\alpha a^* \ , \ \ \bar{v}_\alpha \ &= \ \varepsilon_{\alpha\beta}\chi^*_\beta a \ , \ r \ = \ \chi^+\chi \ + \ a^*a \ . \end{aligned} \tag{9}$$

The operators x_i, b are even generators and $v_\alpha, \bar{v}_\alpha$ the odd ones of the superalgebra $su(2|1)$; r is the central element extending it to the $u(2|1)$ superalgebra.

The $u(2|1)$ superalgebra is realized in the space of $s\mathcal{H}_k$ as the adjoint Poisson-bracket superalgebra $\{X_i, B, V_\alpha, \bar{V}_\alpha, R\}$:

$$X_i\Psi \ = \ [x_i, \Psi] \ , \ B\Psi \ = \ [b, \Psi] \ , \ R\Psi \ = \ [r, \Psi]$$

$$V_\alpha\Psi \ = \ [v_\alpha, \Psi] \ , \ \bar{V}_\alpha\Psi \ = \ [\bar{v}_\alpha, \Psi] \ . \tag{10}$$

Obviously, the space $s\mathcal{H}_k$ is an invariant space with respect to (11). This action can be extended to the superalgebra $\mathcal{B} = \mathcal{U}(u(2|1)) = \mathcal{B}^0 \oplus \mathcal{B}^1$ (the enveloping algebra of the superalgebra $\mathcal{U}(u(2|1))$. Since the function $r = \chi^+\chi + \ a^*a$ is an invariant function with respect to this action, all functions (fields) Φ can be factorized by the relation $r = const.$

The spinor space $\mathcal{S}_k$ is invariant with respect to the action of the even Poisson bracket subalgebra corresponding to $\mathcal{A} := \mathcal{B}^0$. The free Dirac operator is an operator in $\mathcal{S}_k$ which has the following form in terms of $u(2|1)$ generators

$$D_0 \ = \ \frac{1}{2}\varepsilon_{\alpha\beta}(V_\alpha V_\beta \ + \ \bar{V}_\alpha \bar{V}_\beta) \ = \ \varepsilon_{\alpha\beta}(\chi_\beta \partial_{\chi^*_\alpha} a^* \partial_a \ + \ \chi^*_\alpha \partial_{\chi_\beta} a \partial_{a^*}) \ . \tag{11}$$

We stress that this Dirac operator already contains a topological gauge field term (the k-monopole magnetic field). Similarly, the chirality operator can be expressed as

$$\Gamma \ = \ B \ - \ R \ = \ a\partial_a \ - \ a^*\partial_{a^*} \ . \tag{12}$$

The chirality operator anticommutes with the Dirac operator: $D_0\Gamma = -\Gamma D_0$, i.e. the Dirac operator is a chiral odd operator mapping $\mathcal{S}^\pm_k \to \mathcal{S}^\mp_k$. The free spinor field action is defined by

$$S[\Psi, \Psi^*] \ = \ \int d\mu \bar{\Psi} D_0 \Psi \ . \tag{13}$$

Now we shall describe a noncommutative version of the free spinor model. We shall go beyond matrix models starting from the infinite dimensional superalgebra $\mathcal{B} = \mathcal{U}(u(2|1)) = \mathcal{B}^0 \oplus \mathcal{B}^1$ and its even subalgebra $\mathcal{A} := \mathcal{B}^0$. Only the evaluation of the action will be performed at some finite level.

We shall work within the canonical $\pi\mathcal{B}$ realization of the superalgebra $\mathcal{B}$: we insert graded commuting variables a, a^*, χ_α, χ^*_α, $\alpha = 1, 2$ by annihilation and creation operators satisfying graded commutation relations

$$[\chi_\alpha, \chi^*_\beta] \; = \; \delta_{\alpha,\beta} \; , \; [a, a^*] \; = \; 1 \tag{14}$$

(all other elementary commutators vanish). They act in the Fock (super)-space

$$s\mathcal{F} \; = \; \{|n, \nu\rangle \; = \; \frac{1}{\sqrt{n!}}\chi^{*n} a^{*\nu}|0\rangle\} \; = \; s\mathcal{F}^0 \; \oplus s\mathcal{F}^1 \; , \tag{15}$$

possesses a natural grading with respect to the fermion occupation number ν (in eq. (16) n is a two-component multiindex, $\nu = 0, 1$, and $|0\rangle$ is the corresponding normalized vacuum state: $\chi_\alpha|0\rangle = a|0\rangle = 0$). In what follows we shall use the notation $\mathcal{F} = s\mathcal{F}^0$, and we shall simply write $\mathcal{B}$ and $\mathcal{A}$ instead of $\pi\mathcal{B}$ and $\pi\mathcal{A}$.

The generators satisfying in $s\mathcal{F}$ the $\mathcal{U}(u(2|1))$ graded commutation relations (10) are given by eqs. (9) (commuting parameters are replaceded by annihilation and creation operators and graded Poisson brackets by corresponding graded commutators). The Fock subspace

$$s\mathcal{F}_N \; = \; \{|n, \nu\rangle \; , \; |n| + \nu = N\} \; = \; s\mathcal{F}^0_N \; \oplus s\mathcal{F}^1_N \; , \tag{16}$$

is the carrier space of the unitary irreducible representation π'_N of $u(2|1)$ superalgebra. We shall use the notation $\mathcal{F}_N = s\mathcal{F}^0_N$.

We define the space $s\mathcal{H}_k$ of superfields with the winding number $2k$ as the set of operators in the Fock space of the form (1) with $2k = |n| - |m| + \mu - \nu$ fixed. Obviously, $\Phi \in s\mathcal{H}_k$ maps any space $s\mathcal{F}_N$ into the space $s\mathcal{F}_M$, $M = N + 2k$. The $u(2|1)$ superalgebra is realized in $s\mathcal{H}_k$ as the adjoint graded commutator superalgebra (10). Any superfield from $s\mathcal{H}_k$ can be expressed as in (2) with coefficients from $\mathcal{H}_k$ (however now expressed in terms of anihilation and creation operators).

By restricting the action of superfields $\Phi \in s\mathcal{H}_k$ from $s\mathcal{F}$ to $s\mathcal{F}_N$ we obtain a space of mappings $s\mathcal{F}_N \to s\mathcal{F}_M$, $M = N + 2k$. We shall denote them by $s\mathcal{H}_k^{J'}$, $k = \frac{1}{2}(M - N)$, $J' = \frac{1}{2}(M + N - 1) = J - \frac{1}{2}$ (these relations among M, N, J, J' and k are assumed in what follows). From the definition of $s\mathcal{H}_k^{J'}$ it follows that on $s\mathcal{H}_k^{J'}$ acts the direct product $\bar{\pi}'_M \otimes \pi'_N$ of the irreducible representations $\bar{\pi}_M$ and π_N of the superalgebra $u(2|1)$.

Note: Representations π'_N of $u(2|1)$ are well known. They corresponds to the atypical $(q, -q)$ representations of $su(2|1)$, in the notation of [20]: quadratic and cubic Casimir operators vanish and they are classified by the value $R = N$ of the central element in $u(2|1)$ superalgebra. However, their products are indecomposable and have quite pathalogical properties, [20]. This causes problems with the identification of auxiliary fields in the noncommutative case.

Such problems do not occur for the coefficient spaces $\mathcal{H}_k$ of operators acting in $\mathcal{F}_N := s\mathcal{F}_N^0$. The subspaces $\mathcal{H}_k^J$ are defined as the spaces of mappings $\mathcal{F}_N \to \mathcal{F}_M$. In $\mathcal{H}_k^J$ we introduce the scalar product

$$(\Phi_1, \Phi_2)_k^J \; = \; \frac{1}{J+1} \mathrm{Tr}(\Phi_1^* \Phi_2) \ . \tag{17}$$

In $\mathcal{H}_k^J$ acts the direct product representation $\bar{\pi}_M \otimes \pi_N$ of the irreducible $SU(2)$ representations $\bar{\pi}_N$ and π_N in $\mathcal{F}_M$ and $\mathcal{F}_N$ respectively. The direct product $\bar{\pi}_M \otimes \pi_N$ possesses a decomposition to irreducible $SU(2)$ representations, and consequently

$$\mathcal{H}_k^J \ = \ \bigoplus_{j=|k|}^{J} V_k^{Jj} \ , \tag{18}$$

where V_k^{Jj} denotes a $SU(2)$ representation space corresponding to a spin j.

For each J and k we constructed in Appendix A an orthormal basis $\{D_{km}^j,\ j = 0, 1, \ldots, |m| \leq j\}$ in $\mathcal{H}_k$. The orthonormal basis $\{D_{km}^{Jj} = c_{kj}^J D_{km}^j$, $j = 0, \ldots, J,\ |m| \leq j\}$ in $\mathcal{H}_k^J$ with respect to the scalar product (17) is determined by rescaling coefficients c_{kj}^J determined in Appendix A.

We define the isometrical projection operator $C_k^J : \mathcal{H}_k \to \mathcal{H}_k^J$ by $C_k^J D_{km}^j = D_{km}^{Jj}$, and by ${C'}_k^J : \mathcal{H}_k^J \to \mathcal{H}_k$ we denote the reversed isometrical embedding. We can extend this construction by taking $K \geq J$ and defining the isometrical projections $C_k^{KJ} : \mathcal{H}_k^K \to \mathcal{H}_k^J$, together with the reversed isometrical embeddings ${C'}_k^{JK} : \mathcal{H}_k^J \hookrightarrow \mathcal{H}_k^K$. It holds

$$C_k^{KJ} \ = \ C_k^J {C'}_k^K \ , \ {C'}_k^{JK} \ = \ C_k^K {C'}_k^J \ . \tag{19}$$

These are the key objects of our approach which give the prescription how are related to each other for various values of J the (matrix) realizations in $\mathcal{H}_k^J$ of a given field from $\mathcal{H}_k$.

Similarly as in the graded commutative case, we define the space $\mathcal{S}_k$ of spinor fields with a topological winding number $2k$ as the odd subspace of the space $s\mathcal{H}_k$. Such spinor fields can be expanded as

$$\Psi \ = \ f(\chi, \chi^*)a \ + \ g(\chi, \chi^*)a^* \ , \tag{20}$$

where $f \in \mathcal{H}_{k+\frac{1}{2}}$ and $g \in \mathcal{H}_{k-\frac{1}{2}}$. The chirality operator $\Gamma : \mathcal{S}_k \to \mathcal{S}_k$ and the charge conjugation $\mathcal{J} : \mathcal{S}_k \to \mathcal{S}_{-k}$ are defined in the same way as in the commutative case.

The subspaces $\mathcal{S}_k^{J'}$ are again given by the restriction: they are odd mappings from $\mathcal{H}_k^{J'}$, i.e. odd mappings $s\mathcal{F}_N \to s\mathcal{F}_M$ with $M = J' + k + \frac{1}{2}$, $N = J' - k + \frac{1}{2}$. In the space $\mathcal{S}_k^J$ we choose the inner product

$$(\Psi_1, \Psi_2)_k^{J'} \;=\; \frac{1}{2J'+1} s\mathrm{Tr}_N(\bar{\Psi}_1 \Psi_2) \ , \tag{21}$$

where $s\mathrm{Tr}_N$ denotes the supertrace in the space of mappings $s\mathcal{F}_N \to s\mathcal{F}_N$.

Now we shall concentrate our attention to the space of mappings $\mathcal{S}_k \to \mathcal{S}_k$. Such mappings have the form

$$\sum A_i B_i^o \Psi \;:=\; \sum A_i \Psi B_i^* \ , \tag{22}$$

where A_i,B_i are elements from $s\mathcal{H}_0$ such that all products A_iB_i are even (B^o means the right multiplication of Ψ by B^*). In the space of mappings $\mathcal{S}_k \to \mathcal{S}_k$ we have a natural grading induced by the chirality operator: the mapping $\sum A_i B_i^o$ is even (odd) if all A_i and B_i are even (odd). Obviously, $\mathcal{S}_k^\pm \to \mathcal{S}_k^\pm$ for the even mappings and $\mathcal{S}_k^\pm \to \mathcal{S}_k^\mp$ for the odd ones.

In the noncommutative case the free Dirac operator is given by the first equation in (11) but with graded Poisson brackets inserted by graded commutators. The resulting expression can be rewritten in the form

$$D_0 \;=\; v_\alpha \bar{v}_\alpha^o \;-\; \bar{v}_\alpha v_\alpha^o \ , \tag{23}$$

which was proposed earlier in [11]. This is the simplest odd mapping commuting with the $su(2)$ generators $X_i = x_i - x_i^o$, $i = 1, 2, 3$.

The spectrum of the free Dirac operator contains:

(i) non-zero modes

$$E_{jk}^{J'\pm} = \pm\sqrt{(j+k+\frac{1}{2})(j-k+\frac{1}{2})} \ , \ j = |k| + \frac{1}{2}, |k| + \frac{3}{2}, \ldots, J \ , \tag{24}$$

(ii) and $2|k|$ zero-modes corresponding to $j = |k| - \frac{1}{2}$ (if $k \neq 0$).

The corresponding eigenfuctions $\{\Psi_{km}^{J'j\pm}\}$, $j = |k| - \frac{1}{2}, |k| + \frac{1}{2}, \ldots, J'$, $|m| \leq j$ are presented in Appendix A. They form an orthonormal basis in $\mathcal{S}_k^{J'}$. Any spinor field from $\mathcal{S}_k^{J'}$ can be expanded as

$$\Psi \ = \ \sum_{j=|k|+1/2}^{J'} \sum_{|m|<j} (a_{km}^{j+} \Psi_{km}^{J'j+} \ + \ a_{km}^{j-} \Psi_{km}^{J'j-}) \ . \tag{25}$$

For a given k and $J = J' + \frac{1}{2}$ the free spinor field action we take as

$$S_k^J[\Psi, \Psi^*] \ = \ (\Psi, D_0 \Psi)_k^J \ . \tag{26}$$

The spinor field space $\mathcal{S}_k$ is obviously an $\mathcal{A}$-bimodule. The algebra $\mathcal{A} \otimes \mathcal{A}^o$ acts in $\mathcal{S}_k$ as left-multiplication by $A \in \mathcal{A}$ and right-multiplication by $B^* \in \mathcal{A}$ (see (22)). The Dirac operator acts in $\mathcal{A} \otimes \mathcal{A}^o$ as a commutator:

$$D_0(AB^o\Psi) \ = \ [D_0, AB^o]\Psi \ + \ AB^o D_0 \Psi \ . \tag{27}$$

Thus, $D_0 \, : \, AB^o \, \rightarrow \, [D_0, AB^o]$. Consequently, the graded Leibniz rule in $\mathcal{A} \otimes \mathcal{A}^o$ is satisfied. Obviously, this changes the grading of the mapping. The charge conjugation $\mathcal{J}$ induces in the space $\mathcal{S}_k \rightarrow \mathcal{S}_k$ an automorphism $AB^o \rightarrow \mathcal{J}^{-1} AB^o \mathcal{J}$. It is easy to see that the Dirac operator corresponds to an $\mathcal{J}$-odd mapping: $\mathcal{J}^{-1} D_0 \mathcal{J} = -D_0$.

3 Gauge symmetry

In the commutative case the gauge transformations of the spinor field are given in the following way

$$\Psi \ \rightarrow \ \omega \Psi \ , \ \Psi^* \ \rightarrow \ \omega^* \Psi^* \ , \tag{28}$$

where ω is an arbitrary unitary element from $\mathcal{H}_0$. To guarantee the gauge invariance one should replace the free Dirac operator D_0 by the full Dirac operator

$$D \ = \ D_0 \ + \ q_o A \ . \tag{29}$$

Here q_o is a dimensionless coupling constant related to the usual coupling constant q and the radius r of the sphere in question as $q_o \ = \ qr$. The compensating gauge field A transforms inhomogenously under gauge transformations

$$A \ \rightarrow \ A \ + \ \omega[D_0, \omega^*] \ . \tag{30}$$

Since, D_0 already contains the topological (k-monopole) gauge field, the field A is a globally defined gauge field. Such gauge fields can be expressed in terms of the real prepotentials $\lambda \in \mathcal{H}_0$ corresponding to the pure gauge field A (exact 1-form) and $\sigma \in \mathcal{H}_0$ corresponding to the dynamical gauge field A (coexact 1-form). The full Dirac operator can be written in terms of $\tilde{\Omega} = UQP_+ + UQ^{-1}P_-$ with $U = \exp(i\lambda)$ and $Q = \exp(q_o\sigma)$. The formula for D can be rewritten as

$$D \ = \ \tilde{\Omega} D_0 \tilde{\Omega}^* \ . \tag{31}$$

The square of the Dirac operator has the form

$$D^2 \ = \ [\Delta(\Omega) + q_o F(\Omega)]P_+ \ + \ [\Delta(\Omega) - q_o F(\Omega)]P_- \ , \tag{32}$$

where $\Delta(\Omega)$ is the Laplace operator on a sphere with the gauge field included, and the field strenght operator $F(\Omega)$ is

$$F(\Omega) \ = \ q_o^{-1}(\chi_\alpha^* \partial_{\chi_\alpha^*} - \chi_\alpha \partial_{\chi_\alpha}) \ + \ \Delta\sigma \ . \tag{33}$$

The differential term takes in $\mathcal{S}_k$ the constant value $q_o^{-1}k$ and it corresponds to the contribution of the k-monopole, the last term is the field strenght generated by the dynamical field σ.

The pure gauge prepotential λ can be gauged away, and there are no restrictions on it. There are limitations on the dynamical prepotential, since σ enters the Schwinger model field action

$$S[\Psi, \Psi^*, \Omega] \ = \ \frac{1}{4}\int d\nu F^2(\Omega) \ + \ \int d\mu \bar{\Psi} D \Psi \ , \tag{34}$$

which is obviously gauge invariant. Expanding σ into spherical functions (denoted as D^j_{0m}):

$$\sigma \ = \ \sum_{j=0}^{\infty} \sum_{|m|\leq j} b^j_m D^j_{0m} \ , \ b^j_{-m} = (-1)^m b^{j*}_m \ , \tag{35}$$

we obtain a contribution to the action proportional to

$$\sum_{j=0}^{\infty} \sum_{|m|\leq j} j^2(j+1)^2 |b^j_m|^2 \ .$$

This is finite provided that

$$b^j_m \ = \ o(j^{-5/2}) \ . \tag{36}$$

In the noncommutative case we identify the group of gauge transformations with the group of unitary operators $\omega \in \mathcal{H}_0$, and we postulate that the spinor fields $\Psi \in \mathcal{S}_k$ transform under gauge transformations as follows

$$\Psi \ \rightarrow \ \tilde{\omega}\Psi \ = \ \omega\omega^o\Psi \ = \ \omega\Psi\omega^* \ . \tag{37}$$

This is an unitary $\mathcal{J}$-invariant mapping in $\mathcal{S}_k$. In order to obtain a gauge invariant action we introduce compensating gauge degrees of freedom. We shall describe them by the operators of the form $\Omega = UQ^{(+)}P_+ + UQ^{(-)}P_-$, where $U \in \mathcal{H}_0$ is an unitary operator describing pure gauge degrees of freedom and $Q^{(\pm)} \in \mathcal{H}_0$ are positive operators corresponding to the dynamical gauge degrees (they will be specified below). To any Ω we assign the mapping $\tilde{\Omega}$ in $\mathcal{S}_k$:

$$\Psi \ \rightarrow \ \tilde{\Omega}\Psi \ = \ \Omega\Omega^o\Psi \ = \ \Omega\Psi\Omega^* \ . \tag{38}$$

Explicitely, acting by $\tilde{\Omega}$ and $\tilde{\Omega}^*$ on $\Psi = fa + ga^* \in \mathcal{S}_k$ we obtain

$$\begin{aligned} \tilde{\Omega}\Psi \ &= \ UQ^{(+)}fQ^{(-)}U^*a \ + \ UQ^{(-)}gQ^{(+)}U^*a^* \ , \\ \tilde{\Omega}^*\Psi \ &= \ Q^{(+)}U^*fUQ^{(-)}a \ + \ Q^{(-)}U^*gUQ^{(+)}a^* \ . \end{aligned} \tag{39}$$

We take the interacting Dirac operator in the form

$$D \ = \ \tilde{\Omega}D_0\tilde{\Omega}^* \ . \tag{40}$$

Postulating that under gauge transformation

$$\Omega \ \rightarrow \ \omega\Omega \ , \tag{41}$$

we enjoy the transformation property: $D\Psi \rightarrow \omega D\Psi\omega^*$.

Formula (40) for the full Dirac operator has the important property that it preserves the index of the Dirac operator. Any zero-mode Ψ_0 of D_0 yields a zero-mode $\Omega^{*-1}\Psi_0$ is of D and vice versa. Nonzero-modes appear in pairs, if $\Psi_E^{(+)}$ is an eigenstate of the full Dirac operator to the eigenvalue $E, E > 0$, then $\Psi_E^{(-)} = \Gamma\Psi_E^{(+)}$ is an eigenstate to the eigenvalue $-E$.

The square of the Dirac operator is an even operator in $\mathcal{S}_k$. Similarly as in the commutative case, the square of the Dirac operator can be written as

$$D^2 \ = \ [\Delta(\Omega) + q_oF(\Omega)]P_+ \ + \ [\Delta(\Omega) - q_oF(\Omega)]P_- \ , \tag{42}$$

where $\Delta(\Omega)$ is the noncommutative analog of a Laplace operator on a sphere with gauge degrees of freedom included, $F(\Omega)$ is the operator coorresponding to the field strenght and q_o is the coupling constant. The topological (monopole) contribution to the field strenght is obtained by putting $\Omega = 1$ in $F(\Omega)$. In $\mathcal{S}_k$ it takes the same value $q_o^{-1}k$ as in the commutative case.

For a given J and k the spinor field action interacting with gauge field has the form

$$\begin{aligned} S_k^J[\Psi, \Psi^*, \Omega] &= \frac{1}{2J'+1} s\mathrm{Tr}_N[\bar{\Psi} D \Psi] \\ + & \frac{1}{J^2+J-k^2} s\mathrm{Tr}_k^{J'}[F^2(\Omega)P_+ - F^2(\Omega)P_-]\} \ , \end{aligned} \tag{43}$$

where $s\mathrm{Tr}_N$ in the first term denotes the supertrace in the space of mappings $s\mathcal{F}_N \rightarrow s\mathcal{F}_N$, and $s\mathrm{Tr}_k^J$ in the second one denotes the supertrace in the space of mappings $\mathcal{S}_k^{J'} \rightarrow \mathcal{S}_k^{J'}$. The factor in the second term guarantees that the pure topological contribution obtained for $\Omega = 1$ is properly normalized to $q_o^{-2}k^2$. The action (46) is gauge invariant. We can gauge U away and fix the gauge by putting $\Omega = Q^{(+)}P_+ + Q^{(-)}P_-$.

In the noncommutative case the dynamical gauge degrees of freedom are described the hermitean operator $\sigma \in \mathcal{H}_0$ possessing an expansion like in (35), however now in terms of a noncommutative analogs of spherical functions: $D^j_{0m} \in \mathcal{H}_0$, $j = 0, 1, \ldots, |m| \leq j$, (see Appendix A). Since, the noncommutative gauge field action approaches in the commutative limit its commutative form we shall assume the asymptotic behaviour (36) for the expansion coefficients too.

We define the operators $Q^{(\pm)}$ in the Fock space $\mathcal{F}$ by their restrictions $Q^{(\pm)}_{N'}$ to subspaces $\mathcal{F}_{N'}$. For fixed J and k in the action appear restrictions with $N' = J' \pm k \pm \frac{1}{2}$. The maximal value of N' for fixed J and arbitrary $|k| \le J$ is $2J$. Therefore, it is enough to define the the operators $Q^{(\pm)}_{N'}$, $N' \le K$, for some fixed $K > 2J$. Namely, we put

$$Q^{(\pm)}_{N'} \;=\; \exp \Sigma^{(\pm)}_{KN'} \; , \;\; N' \le K \; , \tag{44}$$

where the operators

$$\Sigma^{(\pm)}_{KN'} \;=\; (\ln \circ {C'}^{KN'}_0 \circ \exp \circ C^{KN'}_0)(\pm e_o \sigma) \; , \tag{45}$$

are defined with the help of isometrical mappings $C^{KN'}_0 : \mathcal{H}^K_k \to \mathcal{H}^{N'}_k$ and ${C'}^{KN'}_0 : \mathcal{H}^{N'j}_k \hookrightarrow \mathcal{H}^{Kj}_k$ introduced in Appendix A (e_o is a dimensionless parameter specified later).

This definition guarantees that $\sigma_{KN'} = C^{KN'}_0 \sigma$ posesess an expansion in $\mathcal{F}_{N'}$

$$\sigma_{KN'} \;=\; C^{KN'}_0 \sigma \;=\; \sum_{j=0}^{N'} \sum_{|m| \le j} b^j_m D^{N'j}_{0m} \; , \tag{46}$$

with the truncated subset of coefficients $\{b^j_m, \; j = 0, 1, \ldots, K, \; |m| \le j\}$. Thus, the untruncated modes have the intensities *independent* on N'. The field σ is for us a primary object, whereas $Q^{(\pm)}$ are secondary ones.

4 Quantization and fermionic determinant

We quantize the Schwinger model within the path integral approach. First we describe the quantization in the commutative case (for details see [14]), and then its noncommutative version.

The quantum expectation values of the field functionals $P[\Xi]$ (where $\Xi = \{\Psi, \Psi^*, \sigma, \lambda\}$ denotes the collection of all fields in question) are defined by the formula

$$\langle P[\Xi]\rangle \;=\; Z^{-1}\,\sum_k \int (D\Xi)_k\; P[\Xi]\; e^{-S[\Xi]}\;, \tag{47}$$

with the summation over all topological winding numbers $\kappa = 2k$ and the normalization

$$Z \;=\; \int (D\Xi)_0\; e^{-S[\Xi]}\;. \tag{48}$$

The action $S[\Xi]$ is given in (43). The symbol $(D\Xi)_k = (D\Psi D\Psi^* D\sigma D\lambda)_k$ denotes a formal integration over all field configurations with a given topological number. The field Ψ can be expanded, e.g. into corresponding eigenfuctions of the free Dirac operator

$$\Psi \;=\; \sum_{j=|k|-1/2}^{\infty}\;\sum_{|m|\le j}\,(a^{j+}_{km}\Psi^{j+}_{km} \;+\; a^{j-}_{km}\Psi^{j-}_{km}) \tag{49}$$

with Grassmannian coefficients $a^{j\pm}_{km}$ (for $j = |k| - 1/2$ the expansion contains only one component corresponding to the zero modes). Then $D\Psi_k \sim \prod da^{j\pm}_{km}$ means a formal infinite dimensional Berezin integration. The symbol $D\Psi^*_k$ is defined analogously.

If the field functional $P[\Xi]$ is gauge invariant, then the integrand in (47) does not depends on a pure gauge prepotential λ. A gauge fixing condition in (47) should be imposed, or alternatively, one can factorize out a pure gauge prepotential λ, and take in (47) only the relevant fields $\Xi = \{\Psi, \Psi^*, \sigma\}$. Then $(D\Xi)_k = (D\Psi D\Psi^* D\sigma)_k$. Expanding σ into spherical fuctions (as in (35)) we introduce a formal infinite dimensional integration $D\sigma \sim \prod db^j_m$ (with $b^j_{-m} = (-1)^m b^j_m$).

Thus, the expectation values are defined only formally, and some regularization procedure is needed. In [14] were calculated in this way various non-perturbative quantities. If the functional $P[\Xi]$ in question is a polynomial in spinor fields then the calculation over $(D\Psi D\Psi^*)_k$ can be performed explicitely. This leads to the fermionic determinants and effective actions which are needed for the calculation of fermionic condensates.

The gauge field effective action $S_k^{reg}[\sigma]$ is obtained by integration over $D\Psi D\Psi^*$ in (47) and is given as

$$S_k^{eff}[\sigma] \;=\; S_k[\sigma] \;+\; \Sigma_k^{reg}[\sigma] \;, \tag{50}$$

where $S_k[\sigma]$ is the classical gauge field action and the second term represents the quantum correction defined by the equation

$$C_k^{reg}\mathrm{det}_k D \;=\; \exp(-\Sigma_k^{reg}[\sigma]) \;. \tag{51}$$

Here C_k^{reg} is a field independent regularizating factor and $\mathrm{det}_k D$ denotes the product of non-zero eigenvalues of D in the sector with topological winding number $\kappa = 2k$. The formula derived in [14] reads

$$\Sigma_k^{reg}[\sigma] \;=\; 2\int d\nu(\nabla\sigma)^2 - \pi i|\kappa| - \frac{1}{2}|\kappa|^2 - 2|\kappa|\ln(|\kappa|!) + 2\sum_{n=1}^{|\kappa|} n\ln n \;. \tag{52}$$

The effective action appears in formulas for many important quantities, e.g. the fermionic condensate - the mean value of the field functional $(\Psi,\Psi) = \int d\mu\bar{\Psi}\Psi$. The simplest way how to calculate it consists in adding a "mass" term $m(\Psi,\Psi)$ to the action and using the formula

$$\langle(\Psi,\Psi)\rangle \;=\; -\frac{\partial}{\partial m}\langle e^{-m(\Psi,\Psi)}\rangle|_{m=0} \;. \tag{53}$$

There are two equal contributions $\langle(\Psi,\Psi)\rangle_{\pm 1/2}$ appearing in the numerator in (47) for $k = \pm 1/2$.

In the noncommutative case we shall start from the general formula (47), however there are important differences. We fix J and restrict the admissible range of k to $|k| \leq J$. The symbol $(D\Xi)^J_k = (D\Psi D\Psi^* D\sigma)^J_k$ denotes an integration over all field configurations with a given topological number:

(i) The spinor fields $\Psi \in \mathcal{S}^{J'}_k$ and $\Psi^* \in \mathcal{S}^{J'}_{-k}$ can be expanded into corresponding eigenfuctions of the free Dirac operator with independent Grassmann coefficients $a^{\pm j}_m$, $a^{*\pm j}_m$. The fermionic part of the integration measure $(D\Xi)^J_k$ denotes the finite Berezin integral

$$(D\Psi D\Psi^*)^J_k \;=\; \prod_{|m|\leq|k|-1/2} \frac{da^0_m}{\sqrt{J}} \prod_{j=|k|+1/2}^{J'} \prod_{|m|\leq j} \frac{da^{\pm j}_m}{\sqrt{J}} \frac{da^{*\pm j}_m}{\sqrt{J}} \;, \tag{54}$$

(the first product comes from the contributions of over zero-modes and is present only for $k \neq 0$).

(ii) We restrict σ to $\mathcal{F}_K$ with some fixed $K > 2J$. Expanding σ as in (46) with expansion coefficients $b^j_m = (-1)^m b^j_{-m}$ we put

$$(D\sigma)^J_k \;=\; \prod_{j=0}^{K} db^j_0 \prod_{m=1}^{j} db^j_m \;. \tag{55}$$

There are no problems with the gauge fixing. The gauge group is isomorphic to $SU(K+1)$ possessing a finite volume which can be factorized out. The number of all modes, the dimension of the measure $(D\Xi)^J_k$, is finite, and consequently there are no ultraviolet divergencies. This allows to calculate straightforwardly various non-perturbative quantities.

Let the observable $P[\Xi]$ in (47) be independent of fermion fields. We perform a bosonization of the model by integrating out the fermionic degrees of freedom and obtaining an effective action $S^{eff}_{Jk}[\sigma] = S^J_k[\sigma] + \Sigma^J_k[\sigma]$. Here

$S_k^J[\sigma]$ is the classical noncommutative gauge action (the second term in (43)) and $\Sigma_k^J[\sigma]$ is the quantum correction given as

$$\exp(-\Sigma_k^J[\sigma]) \; = \; \frac{\det_k D}{\det D_0} \; = \; \frac{\int (D'\Psi D'\Psi^*)_k^J e^{-(\bar\Psi, D\Psi)_k^J}}{\int (D\Psi D\Psi^*)_0^J e^{-(\bar\Psi, D_0\Psi)_0^J}} \;, \tag{56}$$

where $(D'\Psi D'\Psi^*)_k^J$ denotes the integration over non-zero modes. It is more convenient to take instead of $D = \tilde\Omega D_0 \tilde\Omega^*$ the shifted Dirac operator $\tilde\Omega(D_0 + m)\tilde\Omega^*$, possessing $2|k|$ modes with $E = m$ instead of zero-modes. Then

$$\exp(-\Sigma_k^J[\sigma]) \; = \; \lim_{m\to 0} \frac{1}{m^{2|k|}} \frac{\int (D\Psi D\Psi^*)_k^J e^{-(\bar\Psi, \tilde\Omega(D_0+m)\tilde\Omega^*\Psi)_k^J}}{\int (D\Psi D\Psi^*)_0^J e^{-(\bar\Psi, D_0\Psi)_0^J}} \;. \tag{57}$$

We rewritte it as a product of two factors. The first one

$$\lim_{m\to 0} \frac{\int (D\Psi D\Psi^*)_k^J e^{-(\bar\Psi, \tilde\Omega(D_0+m)\tilde\Omega\Psi)_k^J}}{\int (D\Psi D\Psi^*)_k^J e^{-(\bar\Psi, (D_0+m)\Psi)_k^J}} \; = \; \det\tilde\Omega \, \det\tilde\Omega^* \tag{58}$$

do not depend on m.

In Appendix B we derived the formula for the rescaling mapping

$$C'^{KN'}_0 \; = \; \exp\{-\sum_{k=0}^{\infty} [(N'+1)^{-2k-1} - (K+1)^{-2k-1}] S_k(\Delta)\} \;, \tag{59}$$

with a known polynomials $S_k(\Delta)$ in Laplace operator $\Delta = X_i^2$, and an analogous formula for $C_0^{KN'}$ differing just by the sign in the exponent. Using them we calculated in Appendix B the commutative limit $J \to \infty$ of the first factor. Taking $K = O(J^\varepsilon)$, $\varepsilon > 1$, and normalizing the constant e_o properly by $e_o^2 = q_o^2(2J+1)^{-1}$ we obtained the leading term in the asymptotic form of (58):

$$\det\tilde\Omega \, \det\tilde\Omega^* \; = \; \exp\{-2q_o^2 \int d\nu (X_i\sigma)^2 \; + \; o(J^{-1})\} \;. \tag{60}$$

The second factor gives with $\kappa = 2k$

$$\lim_{m\to 0} \frac{1}{m^{2|k|}} \frac{\int (D\Psi D\Psi^*)_k^J e^{-(\bar\Psi, (D_0+m)\Psi)_k^J}}{\int (D\Psi D\Psi^*)_0^J e^{-(\bar\Psi, D_0\Psi)_0^J}} \; = \; J^{2|k|^2} \, \frac{\det_k D_0}{\det_0 D_0} \; =$$

$$\exp[\pi i|\kappa| + \frac{1}{2}|\kappa|^2 + 2|\kappa|\ln(|\kappa|!) - 2\sum_{n=1}^{|\kappa|} n\ln n \ + \ o(J^{-1})] \ . \qquad (61)$$

Eqs. (60) and (61) yields in the limit $J \to \infty$ the well-known expression for the effective action (see e.g. [14]).

5 Concluding remarks

In this article we described a noncommutative regularization of the Schwinger model on a noncommutative sphere. We constructed Connes real spectral triple $(\mathcal{A}, \mathcal{H}, D_0)$ supplemented by the the grading Γ and the antilinear isometry $\mathcal{J}$ (identified with the chiralitity operator and charge conjugation). An important new aspect with respect to our previous work [10]-[12] lies in the fact that as the spectral algebra $\mathcal{A}$ we take an infinitedimensional associative algebra - the even part of the superalgebra $\mathcal{B} = \mathcal{U}(u(2|1))$. We have worked within its canonical realization.

The elements $AB^o \in \mathcal{A} \otimes \mathcal{A}^o$ act in the Hilbert space $\mathcal{H}$ of spinor fields as the left multiplications by A and the right one by B^*. Introducing the differentials $dA = [D_0, A]$ and $dB^o = [D_0, B^o]$ we obtain that in our case the condition $[A, dB^o] = [dA, B^o] = 0$, usually required in the noncommutative geometry, is violated.

We gauged the model in a standard way: $\Psi \to \omega\Psi\omega^*$, with ω unitary. The compensating gauge fields A enter the full Dirac operator which can be expressed as $D = D_\alpha\bar{D}_\alpha + D_\alpha^*\bar{D}_\alpha^*$ with $D_\alpha = V_\alpha + q_o A_\alpha$ and $\bar{D}_\alpha = \bar{V}_\alpha + q_o\bar{A}_\alpha$. Such a form of D with general A_α and $\bar{A}_\alpha$ was recently proposed in [17] within supersymmetric extension of the Schwinger model. This formulation contains besides physical fields auxiliary fields. We eliminated them by expressing $A_\alpha = UQ[V_\alpha, U^{-1}Q^{-1}U^{-1}]$ and $\bar{A}_\alpha = UQ^{-1}[\bar{V}_\alpha, QU^{-1}]$ in terms of prepotentials. Moreover, our choice preserves the index of the Dirac operator.

The physical model is determined if we choose the action for the fields in question. The evaluation of the action we performed by restricting the fields to a particular finite dimensional mappings in the Fock space. This required sequences of linear $SU(2)$-invariant embeddings of various representations of $\mathcal{A}$. We evaluated the rescaling coefficients of these embeddings which are necessary for the evaluation of the action.

The action of the resulting noncommutative Schwinger model contains finite number of modes for each field in question. This allows nonperturbatively to quantize the model. As an application we calculated the chiral anomaly and the effective actions. Although we obtained standard results in the commutative limit our interpretation is very different:

- in the commutative case the nontrivial value of the fermionic determinants is due to the singularity of the operator $\exp(-\varepsilon D^2)$ for $\varepsilon \to 0$, and this leads to the chiral symmetry breaking,

- in our approach the chiral anomaly follows straightforwardly from the basic postulates - its appearance is a direct consequence of the fact that the isometric embeddings and exp-log mappings do not commute.

The noncommutative formulation of the Schwinger model leads not only to the expected ultraviolet regularization, but straightforwardly to highly nonperturbative results. It provides a better understanding of the procedure of regularization and renormalization, and the mechanisms of bosonization and chiral symmetry breaking. It would be interesting to extend our investigations in various directions, e.g. to include gravity along lines presented in [3].

Appendix A

Here we shall describe various $SU(2)$ representations in the space $\mathcal{H}_k$. Any operator Φ from this space can be expanded as $\Phi = \sum b^j_m D^j_{km}$ with $m = -j, -j+1, \dots, +j$, $j = |k|, |k|+1, \dots$. The operators D^j_{km} can be constructed as follows:

(i) The operator

$$D^j_{k,-j} \;=\; [(2j+1)!/(j+k)!(j-k)!]^{1/2}\, \hat{\chi}_2^{*j+k} \hat{\chi}_1^{j-k} \tag{62}$$

is the lowest weight with respect to the adjoint action of the operator X_0 because $X_0 D^j_{k,-j} = [x_0, D^j_{k,-j}] = -j D^j_{k,-j}$ and $X_- D^j_{k,-j} = [x_-, D^j_{k,-j}] = 0$ (here $\hat{\chi}_\alpha = \chi_\alpha (\chi + \chi)^{-1/2}$ and $\hat{\chi}^*_\alpha = (\chi + \chi)^{-1/2} \chi^*_\alpha$).

(ii) For a given j all other D^j_{km} can be obtained by a repeated action of X_+,

$$D^j_{km} \;=\; [(j+m)!/(j-m)!(2j)!]^{1/2}\; X_+^{j-m} D^j_{k,-j}\ . \tag{63}$$

The operators D^j_{km}, $|m| \le j$ (for a given j and k) span a representation space V^j_k of the $SU(2)$ representation with spin j.

By definition we take $\{D^j_{km}, j = |k|, |k+1, \dots, |m| \le j\}$ as an orthonormal basis in $\mathcal{H}_k$. Thus,

$$\mathcal{H}_k \;=\; \bigoplus_{j=|k|}^{\infty} V^j_k\ . \tag{64}$$

Restricting the action of superfields $\Phi \in \mathcal{H}_k$ from $s\mathcal{F}$ to $s\mathcal{F}_N$ we obtain a space of mappings $s\mathcal{F}_N \to s\mathcal{F}_M$, $M = N + 2k$. We shall denote it by $\mathcal{H}^J_k$,

$k = \frac{1}{2}(M-N)$, $J = \frac{1}{2}(M+N)$. The expansion (64) is now truncated:

$$\mathcal{H}_k^J \; = \; \bigoplus_{j=|k|}^{J} V_k^{Jj} \; , \tag{65}$$

where V_k^{Jj} denotes the space spanned by the operators (63) restricted to $s\mathcal{F}_N$. If we introduce in $\mathcal{H}_k^J$ the scalar product

$$(\Phi_1, \Phi_2)_k^J \; = \; \frac{1}{J+1} \mathrm{Tr}(\Phi_1^* \Phi_2) \; , \tag{66}$$

then the orthonormal basis in V_k^{Jj} is given as

$$D_{km}^{Jj} \; = \; c_{kj}^J D_{km}^j \; , \; m = -j, \ldots, +j \; , \tag{67}$$

with

$$c_{kj}^J \; = \; [(J+1)(J+k)!(J-k)!/(J+j+1)!(J-j)!]^{1/2} \; . \tag{68}$$

Summing up over $j = |k|, \ldots, J$ one obtains aan orthonormal basis in $\mathcal{H}_k^J$.

Taking $K \geq J$ we define the isometrical projection operator $C_k^{KJ} : \mathcal{H}_k^K \to \mathcal{H}_k^J$ by $D_{km}^{Kj} = C_k^{KJ} D_{km}^{Jj}$, and by ${C'}_k^{KJ} : \mathcal{H}_k^{Jj} \to \mathcal{H}_k^{Kj}$ we denote the reversed isometrical embedding. The operators C_k^{KJ} and ${C'}_k^{KJ}$ are both diagonal with the eigenvalues c_{kj}^K / c_{kj}^J and c_{kj}^J / c_{kj}^K respectively, as follows from the equation

$$c_{kj}^J D_{km}^{Kj} \; = \; c_{kj}^K D_{km}^{Jj} \; , \; j \leq J \leq K \; . \tag{69}$$

Now we determine the spectrum of the free Dirac operator. Since, $[X_0, D_0] = 0$, we classify the eigenstates of D_0 according to the eigenvalues values of X_0. The spinor field

$$\Psi \; = \; \alpha \chi_1^{*m} \chi_2^{n-1} a \; + \; \beta \chi_1^{*m-1} \chi_2^n a^* \tag{70}$$

is the lowest weight of X_0 because, $X_0\Psi = [x_0, \Psi] = (m+n-1)\Psi$ and $X_-\Psi = [x_-, \Psi] = 0$. Moreover,

$$D_0\Psi \;=\; \alpha m \chi_1^{*m-1}\chi_2^n a^* \;+\; \beta n \chi_1^{*m}\chi_2^{n-1} a \;. \tag{71}$$

This is an eigenstate to the value $E = \pm\sqrt{mn}$ provided that $\alpha\sqrt{m} = \beta\sqrt{n}$. The spinor field lowest weight eigenstates are

$$\Psi^{j\pm}_{k,-j} \;=\; c(\sqrt{n}\chi_1^{*m}\chi_2^{n-1} a \;\pm\; \sqrt{m}\chi_1^{*m-1}\chi_2^n a^*) \;, \tag{72}$$

where $j = \frac{1}{2}(m+n-1)$, $k = \frac{1}{2}(m-n)$ (if $k \neq 0$ we obtain for $m=0$ or $n=0$ zero modes corresponding to the first or second term in (74) respectively). The remaining eigenstates of D_0 to the same eigenvalue are given by the standard formula

$$\Psi^{j\pm}_{km} \;=\; [(j+m)!/(j-m)!(2j)!]^{1/2} \;\; X_+^{j-m}\Psi^{j\pm}_{k,-j} \;. \tag{73}$$

The admissible values of j then are $j = |k|, |k|+1, \ldots J$. The normalization constant c in (72) is specified if we restrict the space of spinor fields $\mathcal{S}_k$ to the space $\mathcal{S}_k^J$ with the inner product:

$$(\Psi_1, \Psi_2)_k^J \;=\; \frac{1}{2J+1}\, s\mathrm{Tr}(\bar{\Psi}_1\Psi_2) \;. \tag{74}$$

Appendix B

From the formula (68) for the rescaling coefficitients it follows that

$$c_{0j}^N \;=\; \exp\left[\frac{1}{2}\sum_{n=1}^{j}\ln\frac{1-n/(N+1)}{1+n/(N+1)}\right] \;=$$

$$\exp\left[-\sum_{k=0}^{\infty}(N+1)^{-2k-1}S_k(j(j+1))\right] \;, \tag{75}$$

where $S_k(t) = [(2k+1)(2k+2)]^{-1}t^k + \ldots$ are polynomials defined by

$$S_k(j(j+1)) = \frac{1}{2k+1}\sum_{n=1}^{j} n^{2k+1}\ .$$

The eigenvalues of the operator ${C'}_0^{KN'}$ can be expressed as

$$c_{0j}^{N'}/c_{0j}^{K} \ = \ \exp\{-\sum_{k=0}^{\infty}[(N'+1)^{-2k-1} - (K+1)^{-2k-1}]S_k(j(j+1))\}\ . \qquad (76)$$

The expansion in (75) is convergent provided $j \le N' \le K$. The eigenvalues of the operator $C_0^{KN'}$ differs just by the sign in front of the sum in exponent.

Replacing $j(j+1)$ by the (noncommutative) Laplace operator on a sphere $\Delta = X_i^2$ one obtains directly from (76) the formulas for the operators ${C'}_0^{KN}$ and $C_0^{KN'}$. This allows to derive the leading term formula

$$\begin{aligned} Q_{N'}^{(\pm)} \ &= \ {C'}_0^{KN'}\exp\{C_0^{KN'}(\pm e_o\sigma)\} \ = \\ &[1 - \frac{K-N'}{2(K+1)(N'+1)}\Delta + \ldots]\exp[\pm e_o(1 + \frac{K-N'}{2(K+1)(N'+1)}\Delta + \ldots)\sigma] \\ &= \ \exp[\pm e_o\sigma - e_o^2\frac{K-N'}{2(K+1)(N'+1)}(X_i\sigma)^2 + o(K^{-2}, N'^{-2})]\ . \qquad (77)\end{aligned}$$

Using the well known formula $\det A\otimes B = (\det A)^M(\det B)^N$ (valid for matrices $A \in \mathrm{Mat}(N\otimes N)$ and $B \in \mathrm{Mat}(M\otimes M)$) and the definition of the operators $\tilde\Omega : \mathcal{S}_k^{J'} \to \mathcal{S}_k^{J'}$ it can be easily seen that in the gauge $U = 1$ we have

$$\begin{aligned}\det\tilde\Omega \ &= \ \det[\Omega\Omega^o] \ = \\ &(\det{}_{M-1}Q^{(+)})^{N+1}(\det{}_N Q^{(-)})^M(\det{}_M Q^{(-)})^N(\det{}_{N-1}Q^{(+)})^{M+1}\ , \qquad (78)\end{aligned}$$

where $N = J' - k + \frac{1}{2}$ and $M = J' + k + \frac{1}{2}$ as usually. Inserting here (77) we obtain

$$\begin{aligned}\ln\det\tilde{\Omega} \;=\;& e_o[(N+1)\mathrm{Tr}_{M-1}\sigma - M\mathrm{Tr}_N\sigma - N\mathrm{Tr}_M\sigma(M+1)\mathrm{Tr}_{N-1}\sigma] \\ & -\frac{e_o^2}{2(K+1)}\left[\frac{(N+1)(K-M+1)}{M}\mathrm{Tr}_{M-1}(\sigma)^2 + \frac{M(K-N)}{N+1}\mathrm{Tr}_N(\sigma)^2 + \right. \\ & \left. \frac{N(K-M)}{M+1}\mathrm{Tr}_M(\sigma)^2 + \frac{(M+1)(K-N+1)}{N}\mathrm{Tr}_{N-1}(\sigma)^2 + \right] \qquad (79)\end{aligned}$$

Taking $K = O(J^\varepsilon)$, $\varepsilon > 1$, normalizing the constant e_o by $e_o^2 = q_o^2(2J+1)^{-1}$ and using relations

$$\frac{1}{N'}\mathrm{Tr}_{N'}\sigma \;\to\; \int d\nu\sigma \quad , \quad \frac{1}{N'}\mathrm{Tr}_{N'}(\nabla\sigma)^2 \;\to\; \int d\nu(\nabla\sigma)^2 \ ,$$

(which are valid for $N' \to \infty$ due to the asymptotic behaviour (36)) we obtain in the limit $J \to \infty$ the asymptotic formula

$$\ln\det\tilde{\Omega} \;=\; -q_o^2\int d\nu \; (X_i\sigma)^2 \;+\; o(J^{-1}) \ , \qquad (80)$$

leading sraightforwardly to eq. (60).

For a given J and $k \neq 0$ the free Dirac operator possesses non-zero modes $E_{jk}^{J\pm} = \pm\sqrt{(j+\frac{1}{2})^2 - k^2}$, $j = |k| + \frac{1}{2}, |k| + \frac{3}{2}, \ldots, J$, with the multiplicity $2j+1$. Putting $j' = j + \frac{1}{2}$ and $J' = J + \frac{1}{2}$ we obtain

$$J^{2|k|^2}\ \frac{\det_k D_0}{\det_0 D_0} \ = (-1)^{2k} J^{2|k|^2}\ \frac{\prod_{j'=|k|+1}^{J'}(j'^2 - |k|^2)^{2j'}}{\prod_{j'=1}^{J'} j'^{4j'}} \ =$$

$$(-1)^{2k}\prod_{m=1}^{|k|}\left(\frac{1+m/J'}{1+(m-|k|)/J'}\right)^{2J+2m}\ \prod_{n=1}^{2|k|} n^{2|k|-2n}\ \frac{J^{2|k|^2}}{\prod_{n=1}^{|k|}(J'+n)^{2|k|}} \ . \qquad (81)$$

Denoting $\kappa = 2k$ we obtain in the limit $J \to \infty$ desired eq. (61).

References

[1] A. Connes, Publ. IHES *62* (1986) 257.

[2] A. Connes, Geometrie Noncommutative (Inter Editions, Paris 1990).

[3] A. Connes, *Gravity coupled with matter and the foundation of non commutative geometry*, hep-th/9603053.

[4] M. Dubois-Violette, C. R. Acad. Sci. Paris *307*, Ser. I (1988) 403.

[5] M. Dubois-Violette, R. Kerner and J. Madore, Journ. Math. Phys. *31* (1990) 316.

[6] F. A. Berezin, Commun. Math. Phys. *40* (1975) 153.

[7] J. Hoppe, Elem. Part. Res. J. *80* (1989) 145 .

[8] J. Madore, Journ. Math. Phys. *32* (1991) 332.

[9] H. Grosse and P. Prešnajder, Lett. Math. Phys. *28* (1993) 239.

[10] H. Grosse, C. Klimčík and P. Prešnajder, Int. Journ. Theor. Phys. *35* (1996) 231.

[11] H. Grosse, C. Klimčík and P. Prešnajder, Commun. Math. Phys. *178* (1996) 507.

[12] H. Grosse, C. Klimčík and P. Prešnajder, Commun. Math. Phys. *185* (1997) 155.

[13] H. Grosse and P. Prešnajder, Lett. Math. Phys. *33* (1995) 171.

[14] C. Jayewardena, Helv. Phys. Acta *61* (1988) 638.

[15] H. Grosse and J. Madore, Phys. Lett. *B283* (1992) 218.

[16] H. Grosse, C. Klimčík and P. Prešnajder, *Finite gauge model on truncated sphere*, Proc. of Schladming School, p. 279, Lect. Notes in Phys. (Springer-Verlag Berlin 1996).

[17] C. Klimčík, *Gauge theories on the noncommutative sphere*, preprint IHES/P/97/77, hep-th/9710153.

[18] E. Hawkins, *Quantization of equivariant vector bundles*, preprint Pensylvania State Univ. CGPG 97-98/8-1, q-alg/9708030.

[19] F. A. Berezin, Introduction to Superanalysis, (Reidel, Dordrecht 1987).

[20] M. Scheunert, W. Nahm and V. Rittenberg, Journ. Math. Phys. *18* (1977) 156.